

ARTICHOKE (*CYNARA SCOLYMUS* L.): A REVIEW OF ITS HEALTH-PROMOTING PROPERTIES

A. Di Napoli^{1,2}, F. Germani¹, S. Domingues Da Silva², L. Senatori², F. Parisi², P. Zucchetti¹

¹ Istituto Italiano di Permacultura, Scagnello, Cuneo, Italy

² Diennea Prolife S.r.l., Borgo San Lorenzo, Florence, Italy

E-mail: agnesedinapoli@outlook.com. ORCID: 0000-0001-6807-2439

Doi: 10.36118/pharmadvances.2023.55

SUMMARY

Cynara scolymus L., called artichoke or globe artichoke, is a perennial herbaceous plant cultivated worldwide. This plant is a common component of the Mediterranean diet and has been used as a remedy for health conditions since antiquity. The aim of this review is to find the health-promoting properties of artichoke, conducting a literature search in PubMed. The results show that 119 studies describe these effects, and 17 health benefits of artichoke are reported in the scientific literature. Antioxidant activity and effects on the liver and lipid profile are the main health-promoting properties of this plant. We found that artichoke also improves cardiovascular and gastrointestinal health and exerts anticancer, antimetabolic and antiobesity, prebiotic and probiotic, renoprotective and antidiabetic activities. Only one or two research articles reported the positive effects of this plant on the immune system, arthritis, photoaging, the reproductive system, the nervous system, fungal infections and periodontal diseases. The health benefits are mainly exerted by phenolics. In conclusion, this review shows the health-promoting properties of artichoke. The main beneficial effects are antioxidant activity and effects on lipid profile and the liver, which are mainly mediated by phenolics. The results of the scientific articles described in this review and the molecular mechanisms related to the health benefits of artichoke should be confirmed by future experimental studies.

Key words

Artichoke, *Cynara scolymus* L., antioxidant, hepatoprotective, lipid profile.

Impact statement

Artichoke (*Cynara scolymus* L.) has many health benefits, and the main properties are antioxidant activity and effects on the liver and lipid profile.

INTRODUCTION

Cynara scolymus L. is a plant species which belongs to the family Asteraceae (1). This perennial plant, commonly known as artichoke or globe artichoke, is grown worldwide. Artichoke is endemic to the Mediterranean region and has probably been tamed in southern Italy. The Arabs brought it to other parts of the Mediterranean in medieval times (2). Ancient populations did know artichoke for its nutri-

tional and health-promoting properties (3). The scientific name comes from the Latin word 'cinis' and the Greek word 'skolymos', which mean 'ash' and 'cardoon', respectively (4). Artichoke is an herbaceous plant that can reach about 1,80 meters in height. The flower head is globe-shaped with green and violet external bracts. The receptacle is located in the lower part of the artichoke head and the "choke" made up of bristles is found above it. Many blue-purple



Figure 1. Artichoke. A botanical illustration of artichoke. (*Cynara scolymus* by Adriana Morgante Giornetti).

flowers are arranged in the center of the head (**Figure 1**). Artichoke buds are cut before blooming and the edible part includes the receptacle and the inner bracts (5, 6). This plant is a common ingredient of the Mediterranean diet and is widely utilized for health purposes. Artichoke leaves are mainly used in infusions and extracts for their health-promoting properties (7, 8). This plant contains minerals, vitamins, dietary fibers and bioactive compounds, which are responsible for its beneficial effects. Phenolics include: hydroxycinnamic acids, such as chlorogenic acid (**Figure 2**), caffeic acid and cynarine (**Figure 3**); anthocyanidins, such as cyanidin; flavones, such as apigenin and luteolin (**Figure 4**). Triterpenes and sesquiterpene lactones (e.g., cynaropicrin) (**Figure 5**) are also found in artichoke. Finally, this plant contains inulin, which is a fiber with health-promoting properties (9-13).




Figure 2. Chlorogenic acid. A chemical structure image of chlorogenic acid (Retrieved from: <https://pubchem.ncbi.nlm.nih.gov/compound/1794427#section=2D-Structure>. Accessed: Oct 30, 2024).




Figure 3. Cynarine. A chemical structure image of cynarine (Retrieved from: <https://pubchem.ncbi.nlm.nih.gov/compound/5281769#section=2D-Structure>. Accessed: Oct 30, 2024).




Figure 4. Luteolin. A chemical structure image of luteolin (Retrieved from: <https://pubchem.ncbi.nlm.nih.gov/compound/5280445#section=2D-Structure>. Accessed: Oct 30, 2024).




Figure 5. Cynaropicrin. A chemical structure image of cynaropicrin (Retrieved from: <https://pubchem.ncbi.nlm.nih.gov/compound/119093#section=2D-Structure>. Accessed: Oct 30, 2024).

In this review, we search scientific literature to identify the health benefits of artichoke. This plant has been used for treating health conditions since ancient times and is still largely utilized by healthcare and herbal practitioners. Research has focused on its properties and there is increasing interest in the different beneficial effects and the molecular mechanisms involved in these activities. Our work has great significance, as it includes the latest research findings on this topic.

METHODS

We investigated the beneficial effects of artichoke on human health, searching for scientific articles in the PubMed database (<https://pubmed.ncbi.nlm.nih.gov/>). The following keywords were used: 'artichoke', '*Cynara scolymus*', 'artichoke therapeutic effects', '*Cynara scolymus* therapeutic effects', 'artichoke properties', 'artichoke health benefits', 'artichoke health-promoting properties', 'artichoke phytotherapy', '*Cynara scolymus* phytotherapy', 'artichoke disease treatment' and '*Cynara scolymus* disease treatment'. The article types selected in PubMed were clinical trials, randomized controlled trials, books and documents. We included only previous studies relating to the topic of this review and written in English. We excluded reviews, systematic reviews, meta-analyses and articles which show only negative results or adverse effects of artichoke. We chose scientific studies through an initial screening by reading titles and abstracts. In a second moment, we read the whole text to find the proper scientific articles.

RESULTS AND DISCUSSION

The results of literature search indicate that 1374 articles are included in PubMed. We chose 119 studies after article screening and we identified 17 health-promoting properties of artichoke (**Table 1**).

Table 1. Health-promoting properties of artichoke. The table shows the beneficial properties of artichoke and the scientific articles which report these effects.

Health-promoting effects	References
Effects on the liver	Ahmadi <i>et al.</i> , 2019; Ben Salem <i>et al.</i> , 2017b; Ben Salem <i>et al.</i> , 2019; Celepli <i>et al.</i> , 2022a; Celepli <i>et al.</i> , 2022b; Colak <i>et al.</i> , 2016; Deng <i>et al.</i> , 2022; El-Boshy <i>et al.</i> , 2017; El Morsy and Kamel, 2015; Elsayed Elgarawany <i>et al.</i> , 2020; Frigerio <i>et al.</i> , 2021; Gebhardt, 1997; Gebhardt, 2001; Gebhardt, 2002a; Gebhardt, 2002b; Gebhardt and Fausel, 1997; Heidarian and Rafieian-Kopaei, 2013; Kirchoff <i>et al.</i> , 1994; Küçükgergin <i>et al.</i> , 2010; Kurt <i>et al.</i> , 2014; Kwon <i>et al.</i> , 2018; Lee <i>et al.</i> , 2021b; Liao <i>et al.</i> , 2021; Majnooni <i>et al.</i> , 2021; Mehmetçik <i>et al.</i> , 2008; Menghini <i>et al.</i> , 2010; Metwally <i>et al.</i> , 2011; Miccadei <i>et al.</i> , 2008; Nasef <i>et al.</i> , 2022; Panahi <i>et al.</i> , 2018; Qiang <i>et al.</i> , 2012; Rangboo <i>et al.</i> , 2016; Saénz Rodríguez <i>et al.</i> , 2002; Sharaf El-Deen <i>et al.</i> , 2017; Speroni <i>et al.</i> , 2003; Sümer <i>et al.</i> , 2020; Tang <i>et al.</i> , 2017; Wang <i>et al.</i> , 2021; Wauquier <i>et al.</i> , 2021

Effects on lipid profile	Ben Salem <i>et al.</i> , 2017b; Ben Salem <i>et al.</i> , 2019; Ben Salem <i>et al.</i> , 2022a; Bogavac-Stanojevic <i>et al.</i> , 2018; Bundy <i>et al.</i> , 2008; Deng <i>et al.</i> , 2022; Englisch <i>et al.</i> , 2000; Frigerio <i>et al.</i> , 2021; Gebhardt, 1998; Heidarian and Rafieian-Kopaei, 2013; Ibrahim <i>et al.</i> , 2022; Küçükgergin <i>et al.</i> , 2010; Küskü-Kiraz <i>et al.</i> , 2010; Kwon <i>et al.</i> , 2018; Liao <i>et al.</i> , 2021; Majnooni <i>et al.</i> , 2021; Panahi <i>et al.</i> , 2018; Qiang <i>et al.</i> , 2012; Qinna <i>et al.</i> , 2012; Rangboo <i>et al.</i> , 2016; Rondanelli <i>et al.</i> , 2013; Rondanelli <i>et al.</i> , 2014; Rondanelli <i>et al.</i> , 2019; Shimoda <i>et al.</i> , 2003; Tang <i>et al.</i> , 2017; Wauquier <i>et al.</i> , 2021
Effects on the cardiovascular system	Ben Salem <i>et al.</i> , 2022a; Bogavac-Stanojevic <i>et al.</i> , 2018; Crevar-Sakac <i>et al.</i> , 2016; Juzyszyn <i>et al.</i> , 2008; Küçükgergin <i>et al.</i> , 2010; Li <i>et al.</i> , 2004; Lupattelli <i>et al.</i> , 2004; Roghani-Dehkordi and Kamkhah, 2009; Wang <i>et al.</i> , 2021; Xia <i>et al.</i> , 2014; Zapolska-Downar <i>et al.</i> , 2002
Effects on the gastrointestinal system	Bundy <i>et al.</i> , 2004; Emendörfer <i>et al.</i> , 2005; Holtmann <i>et al.</i> , 2003; Ishida <i>et al.</i> , 2010; Marakis <i>et al.</i> , 2002; Nassar <i>et al.</i> , 2013; Sabater <i>et al.</i> , 2019; Verspohl <i>et al.</i> , 2008; Walker <i>et al.</i> , 2001
Antimetabolic and antiobesity activity	Ardalani <i>et al.</i> , 2020; Ben Salem <i>et al.</i> , 2019; Ben Salem <i>et al.</i> , 2022a; Ebrahimi-Mameghani <i>et al.</i> , 2018; Kwon <i>et al.</i> , 2018; Rezazadeh <i>et al.</i> , 2018a; Rezazadeh <i>et al.</i> , 2018b; Rondanelli <i>et al.</i> , 2014; Wauquier <i>et al.</i> , 2021
Anticancer activity	Abdel-Moneim <i>et al.</i> , 2021; Ding <i>et al.</i> , 2021; Islam <i>et al.</i> , 2021; Lepore <i>et al.</i> , 2019; Liu <i>et al.</i> , 2019; Menghini <i>et al.</i> , 2010; Metwally <i>et al.</i> , 2011; Miccadei <i>et al.</i> , 2008; Mileo <i>et al.</i> , 2012; Mileo <i>et al.</i> , 2015; Mileo <i>et al.</i> , 2020; Muti <i>et al.</i> , 2022; Pulito <i>et al.</i> , 2015; Villarini <i>et al.</i> , 2021; Yang <i>et al.</i> , 2022
Probiotic and prebiotic activities	Costabile <i>et al.</i> , 2010; Fissore <i>et al.</i> , 2015; López-Molina <i>et al.</i> , 2005; Riezzo <i>et al.</i> , 2012; Valerio <i>et al.</i> , 2010; Van den Abbeele <i>et al.</i> , 2020; Zeaiter <i>et al.</i> , 2019
Antioxidant activity	Abdel-Moneim <i>et al.</i> , 2021; Ahmadi <i>et al.</i> , 2019; Ben Salem <i>et al.</i> , 2017a; Ben Salem <i>et al.</i> , 2017b; Ben Salem <i>et al.</i> , 2019; Ben Salem <i>et al.</i> , 2022a; Ben Salem <i>et al.</i> , 2022b; Biel <i>et al.</i> , 2020; Bogavac-Stanojevic <i>et al.</i> , 2018; Brown and Rice-Evans, 1998; Carpentieri <i>et al.</i> , 2022; Celepli <i>et al.</i> , 2022a; Celepli <i>et al.</i> , 2022b; Cicek <i>et al.</i> , 2022; Colak <i>et al.</i> , 2016; Crevar-Sakac <i>et al.</i> , 2016; D'Antuono <i>et al.</i> , 2018; Deng <i>et al.</i> , 2022; Durazzo <i>et al.</i> , 2013; El-Boshy <i>et al.</i> , 2017; El Morsy and Kamel, 2015; Elsayed Elgarawany <i>et al.</i> , 2020; Gebhardt, 1997; Gebhardt and Fausel, 1997; Gurel <i>et al.</i> , 2007; Heidarian and Rafieian-Kopaei, 2013; Ibrahim <i>et al.</i> , 2022; Jiménez-Escrig <i>et al.</i> , 2003; Juzyszyn <i>et al.</i> , 2008; Khattab <i>et al.</i> , 2016; Kostić <i>et al.</i> , 2021; Küçükgergin <i>et al.</i> , 2010; Küskü-Kiraz <i>et al.</i> , 2010; Lee <i>et al.</i> , 2021a; Lee <i>et al.</i> , 2021b; Liao <i>et al.</i> , 2021; Lin <i>et al.</i> , 2022; Liu <i>et al.</i> , 2019; Magielse <i>et al.</i> , 2014; Matsumoto <i>et al.</i> , 2021; Mehmetçik <i>et al.</i> , 2008; Menghini <i>et al.</i> , 2010; Metwally <i>et al.</i> , 2011; Miccadei <i>et al.</i> , 2008; Mohammed <i>et al.</i> , 2020; Nasef <i>et al.</i> , 2022; Nassar <i>et al.</i> , 2013; Pérez-García <i>et al.</i> , 2000; Rezazadeh <i>et al.</i> , 2018a; Sarawek <i>et al.</i> , 2008; Skarpanska-Stejnborn <i>et al.</i> , 2008; Speroni <i>et al.</i> , 2003; Takei <i>et al.</i> , 2015; Tang <i>et al.</i> , 2017; Wang <i>et al.</i> , 2021; Xia <i>et al.</i> , 2014; Zapolska-Downar <i>et al.</i> , 2002
Antidiabetic effects	Ben Salem <i>et al.</i> , 2017b; Deng <i>et al.</i> , 2022; Ebrahimi-Mameghani <i>et al.</i> , 2018; Fantini <i>et al.</i> , 2011; Ibrahim <i>et al.</i> , 2022; Kwon <i>et al.</i> , 2018; Rondanelli <i>et al.</i> , 2014
Antiarthritic effects	Masutani <i>et al.</i> , 2016; Wauquier <i>et al.</i> , 2021
Renoprotective activity	Ben Salem <i>et al.</i> , 2022b; El-Boshy <i>et al.</i> , 2017; Khattab <i>et al.</i> , 2016; Sümer <i>et al.</i> , 2020; Wang <i>et al.</i> , 2021
Neuroprotective effects	Cicek <i>et al.</i> , 2022; Ibrahim <i>et al.</i> , 2022
Effects on the reproductive system	Gurel <i>et al.</i> , 2007; Mohammed <i>et al.</i> , 2020

Effects on the immune system	El-Boshy <i>et al.</i> , 2017; Hueza <i>et al.</i> , 2019
Antiphotoaging effects	Takei <i>et al.</i> , 2015; Tanaka <i>et al.</i> , 2013
Antifungal effects	Zhu <i>et al.</i> , 2005
Prevention of periodontal diseases	Hayata <i>et al.</i> , 2019

Effects on the liver

Several research studies examined the effects of artichoke consumption on the liver, performing *in vivo* and *in vitro* experiments. Two studies showed the beneficial effect of artichoke leaf extract (ALE) alone (14) or in combination with metformin or vitamin E (15) in individuals with non-alcoholic fatty liver disease (NAFLD). Artichoke extracts also improve NAFLD in rodents (16, 17). Rangboo and colleagues (18) showed that ALE exerts hepatoprotective activity in a sample of 60 individuals with non-alcoholic steatohepatitis (NASH), while Tang and colleagues (19) found that an artichoke extract has a beneficial effect on alcoholic liver disease (ALD) in mice. Previous studies demonstrated that ALE exerts hepatoprotective effects *in vivo* (20-35), *in vitro* (36) and *ex vivo* (37). Stem and receptacle extracts have this effect in rats (38) and an extract with high phenolic content exerts hepatoprotective and choleric effects in rats (39). Previous scientific articles reported that artichoke has hepatoprotective effects in these laboratory animals (40-42). Qiang and colleagues (43) found that ALE increases bile acid secretion in hamsters. Artichoke extracts exert choleric activity *in vivo* (44, 45) and *in vitro* (46). Previous studies demonstrated that these plant extracts have hepatoprotective (47-49) and anticholestatic (50, 51) effects *in vitro*, which are mainly exerted by phenolics. Gebhardt (52) showed that liver cholesterol synthesis can be inhibited by ALE *in vitro*, and flavones are the bioactive compounds mainly involved in this effect.

Effects on lipid profile

The effects of artichoke consumption on lipid profile are well documented in scientific liter-

ature. ALE intake alone (14) or in association with metformin or vitamin E (15) improves lipid profile in individuals with NAFLD. A study by Rangboo *et al.* (18) showed that ALE consumption is effective in lowering triglycerides, total cholesterol and low-density lipoprotein cholesterol (LDL-C) in a cohort of 60 individuals with NASH. Shimoda and colleagues (53) demonstrated that the beneficial effect of artichoke on lipid profile is exerted by sesquiterpenes. Another study by Bundy *et al.* (54) showed that ALE can be effective in reducing total cholesterol in individuals with hypercholesterolemia. Englisch and colleagues (55) demonstrated that ALE can lower LDL-C and total cholesterol in a cohort of 143 individuals with hyperlipoproteinemia. A previous study by Rondanelli *et al.* (56) found that ALE intake reduces total cholesterol and LDL-C and raises high-density lipoprotein cholesterol (HDL-C) in a sample of 92 patients with hypercholesterolemia, while other scientific articles reported that artichoke extracts can decrease triglycerides, total cholesterol and LDL-C and improve HDL-C in rats (16, 21). Rondanelli and colleagues (57) demonstrated that ALE improves HDL-C and lowers total cholesterol/HDL-C ratio in a cohort of 20 individuals with mild hypercholesterolemia. Another study found that consumption of an artichoke extract ameliorates lipid profile in a sample of 55 overweight individuals with impaired fasting glycemia (58). Two articles reported the beneficial effect of ALE on cholesterol homeostasis, performing *in vitro* (46) and *ex vivo* (37) experiments. Many research studies found the beneficial effect of artichoke extracts on lipid profile in rodents (19, 22, 28, 29, 31, 32, 43, 59, 60-63). A study by Gebhardt (64) showed that ALE is effective

in suppressing hepatic cholesterol synthesis in rats and luteolin is implicated in this activity.

Effects on the cardiovascular system

Artichoke intake may exert beneficial effects on the cardiovascular system. A previous study by Lupattelli *et al.* (65) found that artichoke leaf juice has beneficial effects on endothelial function in a sample of 28 individuals with hyperlipidemia, while Roghani-Dehkordi and Kamkhah (66) demonstrated that artichoke leaf juice is effective in reducing blood pressure in individuals with mild hypertension. Artichoke extracts exert protective effects on the cardiovascular system *in vivo* (29, 59, 60, 67) and *in vitro* (68, 69). A previous scientific article reported the health benefits of artichoke bud extract in a rat model of hypertension (42). Li and colleagues (70) found that ALE can improve expression and function of endothelial nitric oxide synthase (eNOS), performing *in vitro* and *ex vivo* experiments and flavones are involved in this activity. ALE can also inhibit the expression of inducible nitric oxide synthase (iNOS) in vascular smooth muscle cells and the phytochemical compounds mainly implicated in this effect are cynarine and cyanidin (71).

Effects on the gastrointestinal system

The oral intake of artichoke can improve gastrointestinal health in humans and animal models. Previous studies demonstrated the beneficial effect of ALE on the gastrointestinal system in individuals with functional (72) and mild (73) dyspepsia. Other studies showed that ALE is effective in alleviating irritable bowel syndrome (IBS) symptoms in individuals with this condition (74, 75). Nassar and colleagues (76) found that an artichoke head extract can exert antiulcerogenic activity in rats. Cynaropicrin is the bioactive compound which may exert antispasmodic effects on the gastrointestinal tract of guinea pigs (77) and antigastritis activities in rats (78). Verspohl and colleagues (2008) demonstrated the beneficial effect of ALE on IBS in an experiment performed on the ileum of rats and a previous

study found the anti-colitis activity of artichoke pectin in mice (80).

Antimetabolic and antiobesity activity

The beneficial effects of artichoke consumption on metabolic syndrome and obesity are described in recent research articles. ALE intake may be effective in ameliorating metabolic syndrome biomarkers (81-83). Ardalani and colleagues (84) found that ALE can reduce body mass index (BMI) in overweight individuals, while Wauquier and colleagues (37) demonstrated the protective effect of ALE on obesity and metabolic syndrome, performing an *ex vivo* study. A previous study showed the health benefits of an artichoke extract, studying a cohort of 55 overweight individuals with impaired fasting glycemia (58). Recent research articles demonstrated the antiobesity and antimetabolic syndrome activities of ALE in rodents (22, 31, 59).

Anticancer activity

The anticancer activity of artichoke is well reported in the scientific literature. A previous study showed the protective effect of an artichoke extract on pleural mesothelioma in a sample of 18 individuals with asbestos-related benign pleural disease (85). ALE exerts anti-tumor effect against malignant pleural mesothelioma *in vivo* and *in vitro* (86). Liu and colleagues (87) demonstrated the anticancer activity of cynaropicrin in HeLa cells. This bioactive compound found in artichoke may exert an inhibitory effect on thioredoxin reductase and promote oxidative stress, which leads to apoptosis. Previous studies showed the beneficial effects of this plant in a rat model of hepatocellular carcinoma (41) and the potential anticancer effects of artichoke extracts in human hepatocellular carcinoma (36, 49), uterine leiomyoma (88), breast cancer (89-91) and colon cancer (92) cells. Cynaropicrin may exert anticancer activity in anaplastic thyroid cancer (93) and lung carcinoma (94) cells. Yang and colleagues (95) showed the antitumor effects of this bioactive compound against neuro-

blastoma *in vivo* and *in vitro*. A study by Abdel-Moneim *et al.* (96) found the beneficial effect of artichoke extracts on lung cancer in rats, which is exerted mainly through antioxidant, proapoptotic and antiproliferative activities.

Probiotic and prebiotic activities

Artichoke may exert probiotic and prebiotic activities, as demonstrated by human and *in vitro* studies. The oral intake of artichoke fortified with a probiotic (*i.e.*, *Lactobacillus paracasei*) can improve constipation in individuals with this condition (97, 98). Previous research studies reported the prebiotic activity of long-chain inulin from artichoke in a sample of 31 healthy individuals (99) and *in vitro* (100, 101). Fissore and colleagues (102) demonstrated that artichoke fibers (*i.e.*, inulin and pectin of low degree of methylation) have prebiotic effects *in vitro*. A study by Van den Abbeele *et al.* (103) showed that an artichoke extract exerts this activity *in vitro*.

Antioxidant activity

Artichoke is a source of bioactive compounds, and its antioxidant activity has been demonstrated in many research studies involving humans, animal models and *in vitro*. ALE intake exerts antioxidant activity in individuals with metabolic syndrome, lowering oxidized-LDL (ox-LDL) levels (82). Skarpanska-Stejnborn and colleagues (104) showed the antioxidant effect of ALE in a sample of 22 rowers during the training. The antioxidant activity of cynaropicrin from artichoke was demonstrated in HeLa cells (87) and human keratinocytes (105). Sesquiterpene lactones (106) or cynarine and cyanidin (71) from artichoke can inhibit the expression of iNOS *in vitro*. Many previous studies showed that artichoke extracts exert this activity in rodent models of different health conditions (16, 19-29, 32-34, 40-42, 59-62, 67, 76, 96, 107-112). Lee and colleagues (17) found the protective role of ALE against oxidative stress, performing experiments in NAFLD mice and HepG2 cells. Other studies showed the beneficial effect of artichoke

extracts with high phenolic contents on oxidative stress (39, 113-115). Pérez-García and colleagues (116) demonstrated the antioxidant effect of ALE *in vitro*. This artichoke extract is effective in blocking reactive oxygen species (ROS) generation in human leukocytes and the phytochemicals mainly involved are luteolin, caffeic acid, cynarine and chlorogenic acid. Previous studies showed that artichoke extracts exert antioxidant activity *in vitro* (36, 47-49, 68, 69, 117). Sarawek and colleagues (118) found that luteolin from artichoke is effective in inhibiting xanthine oxidase (XO) *in vitro*. Two previous studies showed the antioxidant effects of these plant extracts, performing *in vivo* and *in vitro* experiments (119, 120). Artichoke seeds (121), rhizome (122), buds (123) and pollen (124) exert antioxidant activity and phenolics are mainly involved in this beneficial effect.

Antidiabetic effects

The consumption of artichoke may have anti-diabetic effects, as demonstrated by research studies on humans and animal models. Rondanelli and colleagues (58) found the beneficial effect of an artichoke extract on glucose metabolism in a sample of 55 overweight individuals with impaired fasting glycemia. Artichoke extracts are effective in improving insulin resistance in individuals with metabolic syndrome (81) and rodents (16, 31, 61). Ben Salem and colleagues (21) found the antihyperglycemic activity of ALE in diabetic rats. The hypoglycemic activity of an artichoke flower head extract was demonstrated in obese and normal rats (125). Few studies have been conducted on humans and further studies using large sample sizes are required to confirm these results. Experiments *in vitro* should clarify the molecular mechanisms involved in these activities.

Antiarthritic effects

Two previous studies found the antiarthritic activity of artichoke. Wauquier and colleagues (37) showed that ALE has a beneficial effect on osteoarthritis, performing an *ex vivo* exper-

iment using human articular chondrocytes. Another study demonstrated the beneficial effect of cynaropicrin on the metabolism of cartilage *in vitro* (126). These two scientific articles represent preliminary evidence that artichoke exerts antiarthritic activity and other studies are required to confirm these results.

Renoprotective activity

The renoprotective effect of artichoke intake has been demonstrated in studies involving animal models. Artichoke bud (42), leaf (25, 107), receptacle and stem (38) extracts may have a protective effect on renal function in rats. Khattab and colleagues (110) found that ALE exerts a beneficial effect on kidney function in a rat model of gentamicin nephrotoxicity.

Neuroprotective effects

ALE exerts a neuroprotective effect in rodent models and this effect may be mediated by the antioxidant activity of artichoke (61, 108). Future studies are needed to confirm these results.

Effects on the reproductive system

Previous studies found that ALE may ameliorate gonadal health in rat models through antioxidant effects (109, 112). The results show the potential beneficial effects of ALE on the reproductive system, but other studies are required to confirm these results and describe in detail the molecular mechanisms underlying these effects.

Effects on the immune system

The oral intake of artichoke may have positive effects on the immune system in animal models. A previous study by El-Boshy *et al.* (25) found that ALE is effective in improving the levels of immunostimulatory cytokines in a rat model of cadmium toxicity. Hueza and colleagues (127) showed that an artichoke extract exerts immunomodulatory activity in rats. These two studies provide the first evidence of the effects of artichoke on the immune system. Other studies are needed to

confirm the results and explain the molecular mechanisms.

Antiphotoaging effects

Cynaropicrin has antiphotoaging effect *in vivo* and *in vitro* by regulating the nuclear factor kappa B (NF- κ B) signaling pathway (128). Takei and colleagues (105) showed that this bioactive compound from artichoke may prevent ultraviolet B (UVB)-induced photoaging in human keratinocytes through antioxidant effects. These results should be confirmed by future studies.

Antifungal effects

Only one research study showed the antifungal activity of different artichoke extracts *in vitro* and found that ALE is the most effective. The authors analyzed the effect against *Candida albicans*, *Candida lusitanae*, *Saccharomyces cerevisiae*, *Saccharomyces carlsbergensis*, *Aspergillus niger*, *Penicillium oxalicum*, *Mucor mucedo* and *Cladosporium cucumerinum* (129). Further studies should corroborate these results.

Prevention of periodontal diseases

Cynaropicrin from artichoke may exert a preventive effect on periodontal diseases *in vitro* by modulating the NF- κ B signaling pathway (130). Only one study found this activity and these results should be confirmed.

This review has some limitations. We reported only scientific articles published in PubMed indexed journals, personal criteria were utilized for conducting the literature search and many studies are characterized by small sample sizes or require replication of results.

CONCLUSIONS

In this study, we show the health-promoting properties of artichoke after performing a literature search. The most common beneficial effects of this plant are those on the liver and lipid profile and antioxidant activity. Other health benefits include improved gastrointes-

tinal and cardiovascular health and anticancer, antimetabolic and antiobesity, prebiotic and probiotic, antidiabetic and renoprotective effects. Only a few studies found the beneficial effects of artichoke on the immune system, the reproductive system, the nervous system, arthritis, photoaging, periodontal diseases and fungal infections. Phenolics are the bioactive compounds mainly involved in these properties and ALE is the extract most utilized for these purposes. Many experiments have been conducted on animal models or *in vitro* and research studies involving humans would be helpful to clarify the beneficial effects of artichoke on human health. The health benefits of artichoke are well documented in scientific literature. Further studies should confirm the results of the articles reported in this review and the molecular mechanisms involved in the health-promoting properties of this plant.

ACKNOWLEDGEMENTS

We are very grateful to Maurizio Carturan and Lidia Novelli for their helpful suggestions.

ETHICS

Funding

There was no institutional or private fundings for this article.

Conflict of interests

The Authors declare that they have no conflict of interests.

Availability of data and material

Data and material are available on request from the Authors.

Authors' contributions

ADN, FG, FP and PZ conceptualized and designed the study. ADN conducted the literature search and drafted the article. All Co-authors discussed the findings, critically revised

the article and approved the final version of the manuscript.

Ethical approval

N/A.

REFERENCES

1. IPNI. International Plant Names Index. The Royal Botanic Gardens, Kew, Harvard University Herbaria & Libraries and Australian National Herbarium. Available from: <http://www.ipni.org>, Accessed: April 15, 2023.
2. Pignone D and Sonnante G. Wild artichokes of south Italy: did the story begin here? *Genetic Resources and Crop Evolution*. 2004;51(6):577-80. Available from: https://www.academia.edu/16978561/Wild_artichokes_of_south_Italy_did_the_story_begin_here. Accessed: Nov 11, 2024.
3. Sonnante G, Pignone D, Hammer K. The domestication of artichoke and cardoon: from Roman times to the genomic age. *Ann Bot*. 2007;100(5):1095-100. doi: 10.1093/aob/mcm127.
4. Verotta L, Macchi MP, Venkatasubramanian P. Connecting Indian wisdom and western science: plant usage for nutrition and health. CRC Press, 2015;316-20.
5. Basay S. Study of the structure of artichoke (*Cynara scolymus* L.) flowers. *JAST*. 2022;24(4):913-24. Available from: <https://jast.modares.ac.ir/article-23-49795-en.html>. Accessed: Nov 11, 2024.
6. Grieve MA. *Modern Herbal*. New York: Dover Publications, 1931.
7. Mulinacci N, Prucher D, Peruzzi M, Romani A, Pinelli P, Giaccherini C, et al. Commercial and laboratory extracts from artichoke leaves: estimation of caffeoyl esters and flavonoidic compounds content. *J Pharm Biomed Anal*. 2004;34(2):349-57. doi: 10.1016/S0731-7085(03)00552-1.
8. Pereira C, Barros L, Carvalho AM, Santos-Buelga C, Ferreira IC. Infusions of

- artichoke and milk thistle represent a good source of phenolic acids and flavonoids. *Food Funct.* 2015;6(1):56-62. doi: 10.1039/c4fo00834k.
9. Azzini E, Bugianesi R, Romano F, Di Venere D, Miccadei S, Durazzo A, et al. Absorption and metabolism of bioactive molecules after oral consumption of cooked edible heads of *Cynara scolymus* L. (cultivar Violetto di Provenza) in human subjects: a pilot study. *Br J Nutr.* 2007;97(5):963-69. doi: 10.1017/S0007114507617218.
 10. Ceccarelli N, Curadi M, Picciarelli P, Martelloni L, Sbrana C, Giovannetti M. Globe artichoke as a functional food. *Mediterr J Nutr Metab.* 2010;3:197-201. doi: 10.1007/s12349-010-0021-z.
 11. Lattanzio V, Kroon PA, Linsalata V, Cardinali A. Globe artichoke: A functional food and source of nutraceutical ingredients. *J Funct Foods.* 2009;1(2):131-44. doi: 10.1016/j.jff.2009.01.002.
 12. Panizzi L and Scarpati ML. Constitution of cynarine, the active principle of the artichoke. *Nature* 1954;174:1062-63. doi: 10.1186/1475-2891-3-5.
 13. Rocchetti G, Lucini L, Corrado G, Colla G, Cardarelli M, Pascale S, et al. Phytochemical profile, mineral content, and bioactive compounds in leaves of seed-propagated artichoke hybrid cultivars. *Molecules.* 2020;25(17):3795. doi: 10.3390/molecules25173795.
 14. Panahi Y, Kianpour P, Mohtashami R, Atkin SL, Butler AE, Jafari R, et al. Efficacy of artichoke leaf extract in non-alcoholic fatty liver disease: A pilot double-blind randomized controlled trial. *Phytother Res.* 2018;32(7):1382-87. doi: 10.1002/ptr.6073.
 15. Majnooni MB, Ataee M, Bahrami G, Heydarpour F, Aneva IY, Farzaei MH, et al. The effects of co-administration of artichoke leaf extract supplementation with metformin and vitamin E in patients with nonalcoholic fatty liver disease: A randomized clinical trial. *Phytother Res.* 2021;35(11):6324-34. doi: 10.1002/ptr.7279.
 16. Deng A, Liu F, Tang X, Wang Y, Xie P, Yang Q, et al. Water extract from artichoke ameliorates high-fat diet-induced non-alcoholic fatty liver disease in rats. *BMC Complement Med Ther.* 2022;22(1):308. doi: 10.1186/s12906-022-03794-9.
 17. Lee M, Kim D, Park SJ, Kim KS, Park GD, Kim OK, et al. Artichoke extract directly suppresses inflammation and apoptosis in hepatocytes during the development of non-alcoholic fatty liver disease. *J Med Food.* 2021;24(10):1058-67. doi: 10.1089/jmf.2021.K.0069.
 18. Rangboo V, Noroozi M, Zavoshy R, Rezadoost SA, Mohammadpoorasl A. The effect of artichoke leaf extract on alanine aminotransferase and aspartate aminotransferase in the patients with nonalcoholic steatohepatitis. *Int J Hepatol.* 2016;2016:4030476. doi: 10.1155/2016/4030476.
 19. Tang X, Wei R, Deng A, Lei T. Protective effects of ethanolic extracts from artichoke, an edible herbal medicine, against acute alcohol-induced liver injury in mice. *Nutrients.* 2017;9(9):1000. doi: 10.3390/nu9091000.
 20. Ahmadi A, Heidarian E, Ghatreh-Samani K. Modulatory effects of artichoke (*Cynara scolymus* L.) leaf extract against oxidative stress and hepatic TNF- α gene expression in acute diazinon-induced liver injury in rats. *J Basic Clin Physiol Pharmacol.* 2019;30(5). doi: 10.1515/jbcpp-2018-0180.
 21. Ben Salem M, Ben Abdallah Kolsi R, Dhouibi R, Ksouda K, Charfi S, Yaich M, et al. Protective effects of *Cynara scolymus* leaves extract on metabolic disorders and oxidative stress in alloxan-diabetic rats. *BMC Complement Altern Med.* 2017;17(1):328. doi: 10.1186/s12906-017-1835-8.
 22. Ben Salem M, Ksouda K, Dhouibi R, Charfi S, Turki M, Hammami S, et al.

- LC-MS/MS analysis and hepatoprotective activity of artichoke (*Cynara scolymus* L.) leaves extract against high fat diet-induced obesity in rats. *Biomed Res Int.* 2019;2019:4851279. doi: 10.1155/2019/4851279.
23. Celepli S, Çolak B, Celepli P, Bigat İ, Batur HG, Soysal F, et al. Artichoke for biochemistry, histology, and gene expression in obstructive jaundice. *Rev Assoc Med Bras.* 2022;68(5):647-52. doi: 10.1590/1806-9282.20220001.
 24. Celepli S, Çolak B, Celepli P, Bigat İ, Batur HG, Soysal F, et al. Effects of artichoke leaf extract on hepatic ischemia-reperfusion injury. *Rev Assoc Med Bras.* 2022;68(1):87-93. doi: 10.1590/1806-9282.20210840.
 25. El-Boshy M, Ashshi A, Gaith M, Qusty N, Bokhary T, AlTaweel N, et al. Studies on the protective effect of the artichoke (*Cynara scolymus*) leaf extract against cadmium toxicity-induced oxidative stress, hepatorenal damage, and immunosuppressive and hematological disorders in rats. *Environ Sci Pollut Res Int.* 2017;24(13):12372-83. doi: 10.1007/s11356-017-8876-x.
 26. El Morsy EM, Kamel R. Protective effect of artichoke leaf extract against paracetamol-induced hepatotoxicity in rats. *Pharm Biol.* 2015;53(2):167-73. doi: 10.3109/13880209.2014.913066.
 27. Elsayed Elgarawany G, Abdou AG, Maher Taie D, Motawea SM. Hepatoprotective effect of artichoke leaf extracts in comparison with silymarin on acetaminophen-induced hepatotoxicity in mice. *J Immunoassay Immunochem.* 2020;41(1):84-96. doi: 10.1080/15321819.2019.1692029.
 28. Heidarian E, Rafieian-Kopaei M. Protective effect of artichoke (*Cynara scolymus*) leaf extract against lead toxicity in rat. *Pharm Biol.* 2013;51(9):1104-09. doi: 10.3109/13880209.2013.777931.
 29. Küçükgergin C, Aydın AF, Ozdemirler-Erta G, Mehmetçik G, Koçak-Toker N, Uysal M. Effect of artichoke leaf extract on hepatic and cardiac oxidative stress in rats fed on high cholesterol diet. *Biol Trace Elem Res.* 2010;135(1-3):264-74. doi: 10.1007/s12011-009-8484-9.
 30. Kurt H, Toprak O, Bülbül E. The possible efficacy of artichoke in flucanazole related hepatotoxicity. *Case Reports Hepatol.* 2014;2014:697359. doi: 10.1155/2014/697359.
 31. Kwon EY, Kim SY, Choi MS. Luteolin-enriched artichoke leaf extract alleviates the metabolic syndrome in mice with high-fat diet-induced obesity. *Nutrients.* 2018;10(8):979. doi: 10.3390/nu10080979.
 32. Liao GC, Jhuang JH, Yao HT. Artichoke leaf extract supplementation lowers hepatic oxidative stress and inflammation and increases multidrug resistance-associated protein 2 in mice fed a high-fat and high-cholesterol diet. *Food Funct.* 2021;12(16):7239-49. doi: 10.1039/d1fo90070f.
 33. Mehmetçik G, Ozdemirler G, Koçak-Toker N, Cevikbaş U, Uysal M. Effect of pretreatment with artichoke extract on carbon tetrachloride-induced liver injury and oxidative stress. *Exp Toxicol Pathol.* 2008;60(6):475-80. doi: 10.1016/j.etp.2008.04.014.
 34. Nasef MA, Yousef MI, Ghareeb DA, Augustyniak M, Aboul-Soud MAM, El Wakil A. Hepatoprotective effects of a chemically-characterized extract from artichoke (*Cynara scolymus* L.) against AFB₁-induced toxicity in rats. *Drug Chem Toxicol.* 2023;46(6):1070-82. doi: 10.1080/01480545.2022.2129672.
 35. Sharaf El-Deen SA, Brakat RM, Mohamed ASAD. Artichoke leaf extract protects liver of *Schistosoma mansoni* infected mice through modulation of hepatic stellate cells recruitment. *Exp Parasitol.* 2017;178:51-59. doi: 10.1016/j.exppara.2017.05.005.
 36. Menghini L, Genovese S, Epifano F, Tirillini B, Ferrante C, Leporini L. Antiproliferative,

- protective and antioxidant effects of artichoke, dandelion, turmeric and rosemary extracts and their formulation. *Int J Immunopathol Pharmacol.* 2010;23(2):601-10. doi: 10.1177/039463201002300222.
37. Wauquier F, Boutin-Wittrant L, Viret A, Guilhaudis L, Oulyadi H, Bourafai-Azies A, et al. Metabolic and anti-inflammatory protective properties of human enriched serum following artichoke leaf extract absorption: results from an innovative ex vivo clinical trial. *Nutrients.* 2021;13(8):2653. doi: 10.3390/nu13082653.
 38. Sümer E, Senturk GE, Demirel ÖU, Yesilada E. Comparative biochemical and histopathological evaluations proved that receptacle is the most effective part of *Cynara scolymus* against liver and kidney damages. *J Ethnopharmacol.* 2020;249:112458. doi: 10.1016/j.jep.2019.112458.
 39. Speroni E, Cervellati R, Govoni P, Guizzardi S, Renzulli C, Guerra MC. Efficacy of different *Cynara scolymus* preparations on liver complaints. *J Ethnopharmacol.* 2003;86(2-3):203-11. doi: 10.1016/s0378-8741(03)00076-x.
 40. Colak E, Ustuner MC, Tekin N, Colak E, Burukoglu D, Degirmenci I, et al. The hepatocurative effects of *Cynara scolymus* L. leaf extract on carbon tetrachloride-induced oxidative stress and hepatic injury in rats. *Springerplus.* 2016;5:216. doi: 10.1186/s40064-016-1894-1.
 41. Metwally NS, Kholeif TE, Ghanem KZ, Farrag AR, Ammar NM, Abdel-Hamid AH. The protective effects of fish oil and artichoke on hepatocellular carcinoma in rats. *Eur Rev Med Pharmacol Sci.* 2011;15(12):1429-44. PMID: 22288304.
 42. Wang ZB, Jiang SL, Liu SB, Peng JB, Hu S, Wang X, et al. Metabolomics of artichoke bud extract in spontaneously hypertensive rats. *ACS Omega.* 2021;6(29):18610-22. doi: 10.1021/acsomega.1c01135.
 43. Qiang Z, Lee SO, Ye Z, Wu X, Hendrich S. Artichoke extract lowered plasma cholesterol and increased fecal bile acids in Golden Syrian hamsters. *Phytother Res.* 2012;26(7):1048-52. doi: 10.1002/ptr.3698.
 44. Kirchhoff R, Beckers C, Kirchhoff GM, Trinczek-Gärtner H, Petrowicz O, Reimann HJ. Increase in choleresis by means of artichoke extract. *Phytomedicine.* 1994;1(2):107-15. doi: 10.1016/S0944-7113(11)80027-9.
 45. Saéñz Rodriguez T, García Giménez D, de la Puerta Vázquez R. Choleric activity and biliary elimination of lipids and bile acids induced by an artichoke leaf extract in rats. *Phytomedicine.* 2002;9(8):687-93. doi: 10.1078/094471102321621278.
 46. Frigerio J, Tedesco E, Benetti F, Insolia V, Nicotra G, Mezzasalma V, et al. Anticholesterolemic activity of three vegetal extracts (artichoke, caigua, and fenugreek) and their unique blend. *Front Pharmacol.* 2021;12:726199. doi: 10.3389/fphar.2021.726199.
 47. Gebhardt R. Antioxidative and protective properties of extracts from leaves of the artichoke (*Cynara scolymus* L.) against hydroperoxide-induced oxidative stress in cultured rat hepatocytes. *Toxicol Appl Pharmacol.* 1997;144(2):279-86. doi: 10.1006/taap.1997.8130.
 48. Gebhardt R, Fausel M. Antioxidant and hepatoprotective effects of artichoke extracts and constituents in cultured rat hepatocytes. *Toxicol In Vitro.* 1997;11(5):669-72. doi: 10.1016/s0887-2333(97)00078-7.
 49. Miccadei S, Di Venere D, Cardinali A, Romano F, Durazzo A, Foddai MS, et al. Antioxidative and apoptotic properties of polyphenolic extracts from edible part of artichoke (*Cynara scolymus* L.) on cultured rat hepatocytes and on human hepatoma cells. *Nutr Cancer.* 2008;60(2):276-83. doi: 10.1080/01635580801891583.
 50. Gebhardt R. Anticholestatic activity of flavonoids from artichoke (*Cynara scolymus* L.) and of their metabolites. *Med*

- Sci Monit. 2001;7(Suppl 1):316-20. PMID: 12211745.
51. Gebhardt R. Prevention of tauroolithocholate-induced hepatic bile canalicular distortions by HPLC-characterized extracts of artichoke (*Cynara scolymus*) leaves. *Planta Med.* 2002;68(9):776-9. doi: 10.1055/s-2002-34417.
 52. Gebhardt R. Inhibition of cholesterol biosynthesis in HepG2 cells by artichoke extracts is reinforced by glucosidase pretreatment. *Phytother Res.* 2002;16(4):368-72. doi: 10.1002/ptr.960.
 53. Shimoda H, Ninomiya K, Nishida N, Yoshino T, Morikawa T, Matsuda H, et al. Anti-hyperlipidemic sesquiterpenes and new sesquiterpene glycosides from the leaves of artichoke (*Cynara scolymus* L.): structure requirement and mode of action. *Bioorg Med Chem Lett.* 2003;13(2):223-28. doi: 10.1016/s0960-894x(02)00889-2.
 54. Bundy R, Walker AF, Middleton RW, Wallis C, Simpson HC. Artichoke leaf extract (*Cynara scolymus*) reduces plasma cholesterol in otherwise healthy hypercholesterolemic adults: a randomized, double-blind placebo-controlled trial. *Phytomedicine.* 2008;15(9):668-75. doi: 10.1016/j.phymed.2008.03.001.
 55. Englisch W, Beckers C, Unkauf M, Ruepp M, Zinserling V. Efficacy of artichoke dry extract in patients with hyperlipoproteinemia. *Arzneimittelforschung.* 2000;50(3):260-65. doi: 10.1055/s-0031-1300196.
 56. Rondanelli M, Giacosa A, Opizzi A, Faliva MA, Sala P, Perna S, et al. Beneficial effects of artichoke leaf extract supplementation on increasing HDL-cholesterol in subjects with primary mild hypercholesterolaemia: a double-blind, randomized, placebo-controlled trial. *Int J Food Sci Nutr.* 2013;64(1):7-15. doi: 10.3109/09637486.2012.700920.
 57. Rondanelli M, Castellazzi AM, Riva A, Allegrini P, Faliva MA, Peroni G, et al. Natural killer response and lipo-metabolic profile in adults with low HDL-cholesterol and mild hypercholesterolemia: beneficial effects of artichoke leaf extract supplementation. *Evid Based Complement Alternat Med.* 2019;2019:2069701. doi: 10.1155/2019/2069701.
 58. Rondanelli M, Opizzi A, Faliva M, Sala P, Perna S, Riva A, et al. Metabolic management in overweight subjects with naive impaired fasting glycaemia by means of a highly standardized extract from *Cynara scolymus*: a double-blind, placebo-controlled, randomized clinical trial. *Phytother Res.* 2014;28(1):33-41. doi: 10.1002/ptr.4950.
 59. Ben Salem M, Affes H, Dhouibi R, Charfi S, Turki M, Hammami S, et al. Effect of artichoke (*cynara scolymus*) on cardiac markers, lipid profile and antioxidants levels in tissue of HFD-induced obesity. *Arch Physiol Biochem.* 2022;128(1):184-94. doi: 10.1080/13813455.2019.1670213.
 60. Bogavac-Stanojevic N, Kotur Stevuljevic J, Cerne D, Zupan J, Marc J, Vujic Z, et al. The role of artichoke leaf tincture (*Cynara scolymus*) in the suppression of DNA damage and atherosclerosis in rats fed an atherogenic diet. *Pharm Biol.* 2018;56(1):138-44. doi: 10.1080/13880209.2018.1434549.
 61. Ibrahim EA, Yousef MI, Ghareeb DA, Augustyniak M, Giesy JP, Aboul-Soud MAM, et al. Artichoke leaf extract-mediated neuroprotection against effects of aflatoxin in male rats. *Biomed Res Int.* 2022;2022:4421828. doi: 10.1155/2022/4421828.
 62. Küskü-Kiraz Z, Mehmetçik G, Dogru-Abbasoglu S, Uysal M. Artichoke leaf extract reduces oxidative stress and lipoprotein dyshomeostasis in rats fed on high cholesterol diet. *Phytother Res.* 2010;24(4):565-70. doi: 10.1002/ptr.2985.
 63. Qinna NA, Kamona BS, Alhussainy TM, Taha H, Badwan AA, Matalka KZ. Effects of prickly pear dried leaves, artichoke leaves, turmeric and garlic extracts, and their combinations on preventing dyslipidemia in

- rats. *ISRN Pharmacol.* 2012;2012:167979. doi: 10.5402/2012/167979.
64. Gebhardt R. Inhibition of cholesterol biosynthesis in primary cultured rat hepatocytes by artichoke (*Cynara scolymus* L.) extracts. *J Pharmacol Exp Ther.* 1998;286(3):1122-28. PMID: 9732368.
65. Lupattelli G, Marchesi S, Lombardini R, Roscini AR, Trinca F, Gemelli F, et al. Artichoke juice improves endothelial function in hyperlipemia. *Life Sci.* 2004;76(7):775-82. doi: 10.1016/j.lfs.2004.07.018.
66. Roghani-Dehkordi F, Kamkhah AF. Artichoke leaf juice contains antihypertensive effect in patients with mild hypertension. *J Diet Suppl.* 2009;6(4):328-41. doi: 10.3109/19390210903280207.
67. Crevar-Sakac M, Vujić Z, Kotur-Stevuljević J, Ivanisević J, Jelić-Ivanović Z, Milenković M, et al. Effects of atorvastatin and artichoke leaf tincture on oxidative stress in hypercholesterolemic rats. *Vojnosanit Pregl.* 2016;73(2):178-87. doi: 10.2298/vsp140917148c.
68. Juzyszyn Z, Czerny B, Pawlik A, Drożdżik M. The effect of artichoke (*Cynara scolymus* L.) extract on ROS generation in HUVEC cells. *Phytother Res.* 2008;22(9):1159-61. doi: 10.1002/ptr.2385.
69. Zapolska-Downar D, Zapolski-Downar A, Naruszewicz M, Siennicka A, Krasnodebska B, Kołodziej B. Protective properties of artichoke (*Cynara scolymus*) against oxidative stress induced in cultured endothelial cells and monocytes. *Life Sci.* 2002;71(24):2897-08. doi: 10.1016/s0024-3205(02)02136-7.
70. Li H, Xia N, Brausch I, Yao Y, Förstermann U. Flavonoids from artichoke (*Cynara scolymus* L.) up-regulate endothelial-type nitric-oxide synthase gene expression in human endothelial cells. *J Pharmacol Exp Ther.* 2004;310(3):926-32. doi: 10.1124/jpet.104.066639.
71. Xia N, Pautz A, Wollscheid U, Reifenberg G, Förstermann U, Li H. Artichoke, cynarin and cyanidin downregulate the expression of inducible nitric oxide synthase in human coronary smooth muscle cells. *Molecules.* 2014;19(3):3654-68. doi: 10.3390/molecules19033654.
72. Holtmann G, Adam B, Haag S, Collet W, Grünewald E, Windeck T. Efficacy of artichoke leaf extract in the treatment of patients with functional dyspepsia: a six-week placebo-controlled, double-blind, multicentre trial. *Aliment Pharmacol Ther.* 2003;18(11-12):1099-105. doi: 10.1046/j.1365-2036.2003.01767.x.
73. Marakis G, Walker AF, Middleton RW, Booth JC, Wright J, Pike DJ. Artichoke leaf extract reduces mild dyspepsia in an open study. *Phytomedicine.* 2002;9(8):694-99. doi: 10.1078/094471102321621287.
74. Bundy R, Walker AF, Middleton RW, Marakis G, Booth JC. Artichoke leaf extract reduces symptoms of irritable bowel syndrome and improves quality of life in otherwise healthy volunteers suffering from concomitant dyspepsia: a subset analysis. *J Altern Complement Med* 2004;10(4):667-9. doi: 10.1089/acm.2004.10.667.
75. Walker AF, Middleton RW, Petrowicz O. Artichoke leaf extract reduces symptoms of irritable bowel syndrome in a post-marketing surveillance study. *Phytother Res.* 2001;15(1):58-61. doi: 10.1002/1099-1573(200102)15:1<58::aid-ptr805>3.0.co;2-r.
76. Nassar MI, Mohamed TK, Elshamy AI, El-Toumy SA, Abdel Lateef AM, Farrag AR. Chemical constituents and anti-ulcerogenic potential of the scales of *Cynara scolymus* (artichoke) heads. *J Sci Food Agric.* 2013;93(10):2494-501. doi: 10.1002/jsfa.6065.
77. Emendörfer F, Emendörfer F, Bellato F, Noldin VF, Cechinel-Filho V, Yunes RA, et al. Antispasmodic activity of fractions and cynaropicrin from *Cynara scolymus* on guinea-pig ileum. *Biol Pharm Bull.* 2005;28(5):902-04. doi: 10.1248/bpb.28.902.
78. Ishida K, Kojima R, Tsuboi M, Tsuda Y, Ito M. Effects of artichoke leaf extract

- on acute gastric mucosal injury in rats. *Biol Pharm Bull.* 2010;33(2):223-29. doi: 10.1248/bpb.33.223.
79. Verspohl EJ, Ploch M, Windeck T, Klaes M, Schmidt T, Bauer K. Effect of two artichoke extracts (36_U and 36_EB) on rat ileum (with respect to bowel syndrome) and the peristaltic threshold. *Phytomedicine.* 2008;15(11):1002-09. doi: 10.1016/j.phymed.2008.02.011.
80. Sabater C, Molina-Tijeras JA, Vezza T, Corzo N, Montilla A, Utrilla P. Intestinal anti-inflammatory effects of artichoke pectin and modified pectin fractions in the dextran sulfate sodium model of mice colitis. Artificial neural network modelling of inflammatory markers. *Food Funct.* 2019;10(12):7793-805. doi: 10.1039/c9fo02221j.
81. Ebrahimi-Mameghani M, Asghari-Jafarabadi M, Rezazadeh K. TCF7L2-rs7903146 polymorphism modulates the effect of artichoke leaf extract supplementation on insulin resistance in metabolic syndrome: a randomized, double-blind, placebo-controlled trial. *J Integr Med.* 2018;16(5):329-34. doi: 10.1016/j.joim.2018.05.006.
82. Rezazadeh K, Aliashrafi S, Asghari-Jafarabadi M, Ebrahimi-Mameghani M. Antioxidant response to artichoke leaf extract supplementation in metabolic syndrome: a double-blind placebo-controlled randomized clinical trial. *Clin Nutr.* 2018;37(3):790-96. doi: 10.1016/j.clnu.2017.03.017.
83. Rezazadeh K, Rahmati-Yamchi M, Mohammadnejad L, Ebrahimi-Mameghani M, Delazar A. Effects of artichoke leaf extract supplementation on metabolic parameters in women with metabolic syndrome: influence of TCF7L2-rs7903146 and FTO-rs9939609 polymorphisms. *Phytother Res.* 2018;32(1):84-93. doi: 10.1002/ptr.5951.
84. Ardalani H, Jandaghi P, Meraji A, Hassanpour Moghadam M. The effect of *Cynara scolymus* on blood pressure and BMI in hypertensive patients: a randomized, double-blind, placebo-controlled, clinical trial. *Complement Med Res.* 2020;27(1):40-6. doi: 10.1159/000502280.
85. Muti P, Sacconi A, Pulito C, Orlandi G, Donzelli S, Morrone A, et al. Artichoke phytocomplex modulates serum microRNAs in patients exposed to asbestos: a first step of a phase II clinical trial. *J Exp Clin Cancer Res.* 2022;41(1):255. doi: 10.1186/s13046-022-02455-6.
86. Pulito C, Mori F, Sacconi A, Casadei L, Ferraiuolo M, Valerio MC, et al. *Cynara scolymus* affects malignant pleural mesothelioma by promoting apoptosis and restraining invasion. *Oncotarget.* 2015;6(20):18134-50. doi: 10.18632/oncotarget.4017.
87. Liu T, Zhang J, Han X, Xu J, Wu Y, Fang J. Promotion of HeLa cells apoptosis by cynaropicrin involving inhibition of thioredoxin reductase and induction of oxidative stress. *Free Radic Biol Med.* 2019;135:216-26. doi: 10.1016/j.freeradbiomed.2019.03.014.
88. Islam MS, Greco S, Delli Carpini G, Giannubilo SR, Segars J, Ciavattini A, et al. Hop and artichoke extracts inhibit expression of extracellular matrix components in uterine leiomyoma cells. *F S Sci* 2021;2(4):407-18. doi: 10.1016/j.xfss.2021.09.002.
89. Mileo AM, Di Venere D, Linsalata V, Fraioli R, Miccadei S. Artichoke polyphenols induce apoptosis and decrease the invasive potential of the human breast cancer cell line MDA-MB231. *J Cell Physiol.* 2012;227(9):3301-09. doi: 10.1002/jcp.24029.
90. Mileo AM, Di Venere D, Abbruzzese C, Miccadei S. Long term exposure to polyphenols of artichoke (*Cynara scolymus* L.) exerts induction of senescence driven growth arrest in the MDA-MB231 human breast cancer cell line. *Oxid Med Cell Longev.* 2015;2015:363827. doi: 10.1155/2015/363827.

91. Mileo AM, Di Venere D, Mardente S, Miccadei S. Artichoke polyphenols sensitize human breast cancer cells to chemotherapeutic drugs via a ROS-mediated down-regulation of flap endonuclease 1. *Oxid Med Cell Longev*. 2020;2020:7965435. doi: 10.1155/2020/7965435.
92. Villarini M, Acito M, di Vito R, Vanini S, Dominici L, Fatigoni C, et al. Pro-apoptotic activity of artichoke leaf extracts in human HT-29 and RKO colon cancer cells. *Int J Environ Res Public Health*. 2021;18(8):4166. doi: 10.3390/ijerph18084166.
93. Lepore SM, Maggisano V, Lombardo GE, Maiuolo J, Mollace V, Bulotta S, et al. Antiproliferative effects of cynaropicrin on anaplastic thyroid cancer cells. *Endocr Metab Immune Disord Drug Targets*. 2019;19(1):59-66. doi: 10.2174/1871530318666180928153241.
94. Ding Z, Xi J, Zhong M, Chen F, Zhao H, Zhang B, et al. Cynaropicrin induces cell cycle arrest and apoptosis by inhibiting PKM2 to cause DNA damage and mitochondrial fission in A549 cells. *J Agric Food Chem*. 2021;69(45):13557-67. doi: 10.1021/acs.jafc.1c05394.
95. Yang R, Ma S, Zhuo R, Xu L, Jia S, Yang P, et al. Suppression of endoplasmic reticulum stress-dependent autophagy enhances cynaropicrin-induced apoptosis via attenuation of the P62/Keap1/Nrf2 pathways in neuroblastoma. *Front Pharmacol*. 2022;13:977622. doi: 10.3389/fphar.2022.977622.
96. Abdel-Moneim A, Ahmed OM, Abd El-Twab SM, Zaky MY, Bakry LN. Prophylactic effects of *Cynara scolymus* L. leaf and flower hydroethanolic extracts against diethylnitrosamine/acetylaminoflourene-induced lung cancer in Wistar rats. *Environ Sci Pollut Res Int*. 2021;28(32):43515-27. doi: 10.1007/s11356-021-13391-x.
97. Riezzo G, Orlando A, D'Attoma B, Guerra V, Valerio F, Lavermicocca P, et al. Randomised clinical trial: efficacy of *Lactobacillus paracasei*-enriched artichokes in the treatment of patients with functional constipation--a double-blind, controlled, crossover study. *Aliment Pharmacol Ther*. 2012;35(4):441-50. doi: 10.1111/j.1365-2036.2011.04970.x.
98. Valerio F, Russo F, de Candia S, Riezzo G, Orlando A, Lonigro SL, et al. Effects of probiotic *Lactobacillus paracasei*-enriched artichokes on constipated patients: a pilot study. *J Clin Gastroenterol*. 2010;44(Suppl 1):S49-53. doi: 10.1097/MCG.0b013e3181d2dca4.
99. Costabile A, Kolida S, Klinder A, Gietl E, B auerlein M, Froberg C, et al. A double-blind, placebo-controlled, crossover study to establish the bifidogenic effect of a very-long-chain inulin extracted from globe artichoke (*Cynara scolymus*) in healthy human subjects. *Br J Nutr*. 2010;104(7):1007-17. doi: 10.1017/S0007114510001571.
100. L pez-Molina D, Navarro-Mart nez MD, Rojas Melgarejo F, Hiner AN, Chazarra S, Rodr guez-L pez JN. Molecular properties and prebiotic effect of inulin obtained from artichoke (*Cynara scolymus* L.). *Phytochemistry*. 2005;66(12):1476-84. doi: 10.1016/j.phytochem.2005.04.003.
101. Zeaiter Z, Regonesi ME, Cavini S, Labra M, Sello G, Di Gennaro P. Extraction and characterization of inulin-type fructans from artichoke wastes and their effect on the growth of intestinal bacteria associated with health. *Biomed Res Int*. 2019;2019:1083952. doi: 10.1155/2019/1083952.
102. Fissore EN, Santo Domingo C, Gerschenon LN, Giannuzzi L. A study of the effect of dietary fiber fractions obtained from artichoke (*Cynara cardunculus* L. var. *scolymus*) on the growth of intestinal bacteria associated with health. *Food Funct*. 2015;6(5):1667-74. doi: 10.1039/c5fo00088b.
103. Van den Abbeele P, Ghyselincx J, Marzorati M, Villar A, Zangara A, Smidt CR,

- et al. In vitro evaluation of prebiotic properties of a commercial artichoke inflorescence extract revealed bifidogenic effects. *Nutrients*. 2020;12(6):1552. doi: 10.3390/nu12061552.
104. Skarpanska-Stejnborn A, Pilaczynska-Szczesniak L, Basta P, Deskur-Smielcka E, Horoszkiewicz-Hassan M. The influence of supplementation with artichoke (*Cynara scolymus* L.) extract on selected redox parameters in rowers. *Int J Sport Nutr Exerc Metab*. 2008;18(3):313-27. doi: 10.1123/ijsnem.18.3.313.
105. Takei K, Hashimoto-Hachiya A, Takahara M, Tsuji G, Nakahara T, Furue M. Cynaropicrin attenuates UVB-induced oxidative stress via the AhR-Nrf2-Nqo1 pathway. *Toxicol Lett*. 2015;234(2):74-80. doi: 10.1016/j.toxlet.2015.02.007.
106. Matsumoto T, Nakashima S, Nakamura S, Hattori Y, Ando T, Matsuda H. Inhibitory effects of cynaropicrin and related sesquiterpene lactones from leaves of artichoke (*Cynara scolymus* L.) on induction of iNOS in RAW264.7 cells and its high-affinity proteins. *J Nat Med*. 2021;75(2):381-92. doi: 10.1007/s11418-020-01479-6.
107. Ben Salem M, Affes H, Dhouibi R, Charfi S, Turki M, Hammami S, et al. Preventive effect of Artichoke (*Cynara scolymus* L.) in kidney dysfunction against high fat-diet induced obesity in rats. *Arch Physiol Biochem*. 2022;128(3):586-92. doi: 10.1080/13813455.2019.1703755.
108. Cicek B, Genc S, Yeni Y, Kuzucu M, Cetin A, Yildirim S, et al. Artichoke (*Cynara scolymus*) methanolic leaf extract alleviates diethylnitrosamine-induced toxicity in BALB/c mouse brain: involvement of oxidative stress and apoptotically related Klotho/PPAR γ signaling. *J Pers Med*. 2022;12(12):2012. doi: 10.3390/jpm12122012.
109. Gurel E, Caner M, Bayraktar L, Yilmazer N, Dogruman H, Demirci C. Effects of artichoke extract supplementation on gonads of cadmium-treated rats. *Biol Trace Elem Res*. 2007;119(1):51-9. doi: 10.1007/s12011-007-0048-2.
110. Khattab HA, Wazzan MA, Al-Ahdab MA. Nephroprotective potential of artichoke leaves extract against gentamicin in rats: antioxidant mechanisms. *Pak J Pharm Sci*. 2016;29(Suppl 5):1775-82. PMID: 28476701.
111. Magielse J, Verlaet A, Breynaert A, Keenoy BM, Apers S, Pieters L, et al. Investigation of the in vivo antioxidative activity of *Cynara scolymus* (artichoke) leaf extract in the streptozotocin-induced diabetic rat. *Mol Nutr Food Res*. 2014;58(1):211-15. doi: 10.1002/mnfr.201300282.
112. Mohammed ET, Radi AM, Aleya L, Abdel-Daim MM. *Cynara scolymus* leaves extract alleviates nandrolone decanoate-induced alterations in testicular function and sperm quality in albino rats. *Environ Sci Pollut Res Int*. 2020;27(5):5009-17. doi: 10.1007/s11356-019-07302-4.
113. Biel W, Witkowicz R, Piątkowska E, Podsiadło C. Proximate composition, minerals and antioxidant activity of artichoke leaf extracts. *Biol Trace Elem Res*. 2020;194(2):589-95. doi: 10.1007/s12011-019-01806-3.
114. Brown JE, Rice-Evans CA. Luteolin-rich artichoke extract protects low density lipoprotein from oxidation in vitro. *Free Radic Res*. 1998;29(3):247-55. doi: 10.1080/10715769800300281.
115. D'Antuono I, Carola A, Sena LM, Linsalata V, Cardinali A, Logrieco AF, et al. Artichoke polyphenols produce skin anti-age effects by improving endothelial cell integrity and functionality. *Molecules*. 2018;23(11):2729. doi: 10.3390/molecules23112729.
116. Pérez-García F, Adzet T, Cañigüeral S. Activity of artichoke leaf extract on reactive oxygen species in human leukocytes. *Free Radic Res*. 2000;33(5):661-65. doi: 10.1080/10715760000301171.

117. Carpentieri S, Augimeri G, Ceramella J, Vivacqua A, Sinicropi MS, Pataro G, et al. Antioxidant and anti-inflammatory effects of extracts from pulsed electric field-treated artichoke by-products in lipopolysaccharide-stimulated human THP-1 macrophages. *Foods*. 2022;11(15):2250. doi: 10.3390/foods11152250.
118. Sarawek S, Feistel B, Pischel I, Butterweck V. Flavonoids of *Cynara scolymus* possess potent xanthinoxidase inhibitory activity in vitro but are devoid of hypouricemic effects in rats after oral application. *Planta Med*. 2008;74(3):221-27. doi: 10.1055/s-2008-1034316.
119. Ben Salem M, Affes H, Athmouni K, Ksouda K, Dhouibi R, Sahnoun Z, et al. Chemicals compositions, antioxidant and anti-inflammatory activity of *Cynara scolymus* leaves extracts, and analysis of major bioactive polyphenols by HPLC. *Evid Based Complement Alternat Med*. 2017;2017:4951937. doi: 10.1155/2017/4951937.
120. Jiménez-Escrig A, Dragsted LO, Daneshvar B, Pulido R, Saura-Calixto F. In vitro antioxidant activities of edible artichoke (*Cynara scolymus* L.) and effect on biomarkers of antioxidants in rats. *J Agric Food Chem*. 2003;51(18):5540-45. doi: 10.1021/jf030047e.
121. Durazzo A, Foddai MS, Temperini A, Azzini E, Venneria E, Lucarini M, et al. Antioxidant properties of seeds from lines of artichoke, cultivated cardoon and wild cardoon. *Antioxidants*. 2013;2(2):52-61. doi: 10.3390/antiox2020052.
122. Lee CL, Liao KC, Chen CC, Lin YA, Wu TY, Jhan YL, et al. Characterization of secondary metabolites from the rhizome of *Cynara scolymus* and their antioxidant properties. *Nat Prod Res*. 2021;35(12):2051-55. doi: 10.1080/14786419.2019.1645664.
123. Lin X, Lu XK, Zhu KH, Jiang XY, Chen JC, Yan PZ, et al. Synchronous extraction, antioxidant activity evaluation, and composition analysis of carbohydrates and polyphenols present in artichoke bud. *Molecules*. 2022;27(24):8962. doi: 10.3390/molecules27248962.
124. Kostić AŽ, Milinčić DD, Nedić N, Gašić UM, Špirović Trifunović B, Vojt D, et al. Phytochemical profile and antioxidant properties of bee-collected artichoke (*Cynara scolymus*) pollen. *Antioxidants*. 2021;10(7):1091. doi: 10.3390/antiox10071091.
125. Fantini N, Colombo G, Giori A, Riva A, Morazzoni P, Bombardelli E, et al. Evidence of glycemia-lowering effect by a *Cynara scolymus* L. extract in normal and obese rats. *Phytother Res*. 2011;25(3):463-66. doi: 10.1002/ptr.3285.
126. Masutani T, Tanaka YT, Kojima H, Tsuboi M, Hara A, Niwa M. Cynaropicrin is dual regulator for both degradation factors and synthesis factors in the cartilage metabolism. *Life Sci*. 2016;158:70-77. doi: 10.1016/j.lfs.2016.06.028.
127. Hueza IM, Gotardo AT, da Silva Mattos MI, Górniak SL. Immunomodulatory effect of *Cynara scolymus* (artichoke) in rats. *Phytother Res*. 2019;33(1):167-73. doi: 10.1002/ptr.6210.
128. Tanaka YT, Tanaka K, Kojima H, Hamada T, Masutani T, Tsuboi M, et al. Cynaropicrin from *Cynara scolymus* L. suppresses photoaging of skin by inhibiting the transcription activity of nuclear factor-kappa B. *Bioorg Med Chem Lett*. 2013;23(2):518-23. doi: 10.1016/j.bmcl.2012.11.034.
129. Zhu XF, Zhang HX, Lo R. Antifungal activity of *Cynara scolymus* L. extracts. *Fito-terapia* 2005;76(1):108-11. doi: 10.1016/j.fitote.2004.10.016.
130. Hayata M, Watanabe N, Kamio N, Tamura M, Nodomi K, Tanaka K, et al. Cynaropicrin from *Cynara scolymus* L. suppresses *Porphyromonas gingivalis* LPS-induced production of inflammatory cytokines in human gingival fibroblasts and RANKL-induced osteoclast differentiation in RAW264.7 cells. *J Nat Med*. 2019;73(1):114-23. doi: 10.1007/s11418-018-1250-6.